

1994 Perry Grass Variety Trial

Brand-Vari * Experiment	4-Jun		7-Jul		1997 7-Aug		10-Sep		Total	1996 Total		1995 Total		3-yr. Total	RelativeYie % Trial Me	
	T	DM/A	T	DM/A	T	DM/A	T	DM/A	T	DM/A	T	DM/A	T	DM/A	T	DM/A
Tall Fescue																
HIMAG*	3.06		1.36		0.41		0.67		5.5		2.48		0.93		8.9	118
Orchardgrass																
WARRIOR	3.23		1.03		0.27		0.45		4.97		1.97		1.1		8.04	107
CONDOR	2.67		1.04		0.25		0.43		4.39		2.01		1.36		7.76	103
POTOMAC	3		0.91		0.18		0.33		4.41		1.78		0.88		7.06	94
Smooth Bromegrass																
LINCOLN	2.57		0.96		0.22		0.34		4.08		1.79		1.15		7.02	93
BRO-MAX	2.92		0.83		0.29		0.26		4.29		1.54		0.65		6.48	86
Mean	2.91		1.02		0.27		0.41		4.61		1.93		1.01		7.54	
5% LSD	0.44		0.26		0.11		0.14		0.7		0.47		0.53		0.93	
CV (%)	10.12		17.19		26.95		22.32		10.07		16.31		34.79		8.15	
MCV (%)	15.25		25.91		40.61		33.63		15.17		24.59		52.43		12.28	

Date of Seeding: April 20, 1994.

Soil Type: Rozetta silt loam.

Herbicides: None.

Fertilization: April 9, June 5-100 lbs. (34-0-0), and June 5 (0-60-250).

Pest Control: None.

Plot Dimensions: 3x16.5'.

ld  
an